| | A | B | C | D | E | F | G | H | I | J | K | L | M | N | O | P | Q | R | S | T | U | V | W | X | Y | Z |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **STATS** | Total vegetation | Ceratophyllum demersum ,Coontail | Elodea canadensis ,Common waterweed | Heteranthera dubia ,Water star-grass | Lemna trisulca ,Forked duckweed | Megalodonta beckii ,Water marigold | Myriophyllum sibericum ,Northern water milfoil | Myriophyllum spicatum ,Eurasian water milfoil | Najas flexilis ,Bushy pondweed | Nitella sp.Nitella | Potamogeton amplifolius Large-leaf pondweed | Potamogeton epihydrus ,Ribbon leaf pondweed | Potamogeton foliosus ,Leafy pondweed | Potamogeton friesii ,Fries's pondweed | Potamogeton nodosus ,Long-leaf pondweed | Potamogeton pusillus,Small pondweed | Potamogeton richardsonii ,Clasping-leaf pondweed | Potamogeton zosteriformis ,Flat-stem pondweed | Ranunculus aquatilis ,Stiff water crowfoot | Sagittaria graminea ,Grass-leaved arrowhead | Vallisneria americana ,Wild celery | | | | |
| 2 | **Total number of points sampled** | **504** | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | **Total number of sites with vegetation** | **230** | | | | | | | | | | | | | | | | | | | | | | | | |
| 4 | **Total number of sites shallower than maximum depth of plants** | **292** | | | | | | | | | | | | | | | | | | | | | | | | |
| 5 | **Frequency of occurrence within vegetated areas (%)** | | 29.56522 | 44.34783 | 13.47826 | 8.26087 | 2.173913 | 6.956522 | 31.73913 | 23.04348 | 0.434783 | 2.173913 | 1.73913 | 4.347826 | 1.73913 | 2.173913 | 15.65217 | 0.869565 | 14.78261 | 4.782609 | 1.73913 | 88.26087 | | | | |
| 6 | **Frequency of occurrence at sites shallower than maximum depth of plants** | **78.8** | 23.28767 | 34.93151 | 10.61644 | 6.506849 | 1.712329 | 5.479452 | 25 | 18.15068 | 0.342466 | 1.712329 | 1.369863 | 3.424658 | 1.369863 | 1.712329 | 12.32877 | 0.684932 | 11.64384 | 3.767123 | 1.369863 | 69.52055 | | | | |
| 7 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 8 | **Relative Frequency (%)** | **100.0** | 9.9 | 14.9 | 4.5 | 2.8 | 0.7 | 2.3 | 10.6 | 7.7 | 0.1 | 0.7 | 0.6 | 1.5 | 0.6 | 0.7 | 5.2 | 0.3 | 5.0 | 1.6 | 0.6 | 29.6 | | | | |
| 9 | Relative Frequency (squared) | 0.146 | 0.010 | 0.022 | 0.002 | 0.001 | 0.000 | 0.001 | 0.011 | 0.006 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.003 | 0.000 | 0.002 | 0.000 | 0.000 | 0.088 | | | | |
| 10 | **Simpson Diversity Index** | **0.85** | | | | | | | | | | | | | | | | | | | | | | | | |
| 11 | **Maximum depth of plants (ft)** | **10** | | | | | | | | | | | | | | | | | | | | | | | | |
| 12 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 13 | **Number of sites with Eurasian water-milfoil** | **73** | | | | | | | | | | | | | | | | | | | | | | | | |
| 14 | **Average Rake Fullness Eurasian water-milfoil** | **1.219178** | | | | | | | | | | | | | | | | | | | | | | | | |
| 15 | **Number of sites with Curly-leaf pondweed** | **#REF!** | | | | | | | | | | | | | | | | | | | | | | | | |
| 16 | **Average Rake Fullness Curly-leaf pondweed** | **#REF!** | | | | | | | | | | | | | | | | | | | | | | | | |
| 17 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 18 | **Number of sites where species found** | | 68 | 102 | 31 | 19 | 5 | 16 | 73 | 53 | 1 | 5 | 4 | 10 | 4 | 5 | 36 | 2 | 34 | 11 | 4 | 203 | | | | |
| 19 | **Number of sites where species found using rake on Pole (P)** | 313 | | | | | | | | | | | | | | | | | | | | | | | | |
| 20 | **Number of sites where species found using rake on Rope (R)** | 0 | | | | | | | | | | | | | | | | | | | | | | | | |
| 21 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 22 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 23 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 24 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 25 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 26 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 27 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 28 | **Number of sites deeper than sample pole (>12 feet deep)** | 192 | | | | | | | | | | | | | | | | | | | | | | | | |
| 29 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 30 | | **Number** | **Percent** | | | | | | | | | | | | | | | | | | | | | | | |
| 31 | **Number of sites with muck** | 112 | 35.9% | | | | | | | | | | | | | | | | | | | | | | | |
| 32 | **Number of sites with sand & gravel** | 117 | 37.5% | | | | | | | | | | | | | | | | | | | | | | | |
| 33 | **Number of sites with rock** | 83 | 26.6% | | | | | | | | | | | | | | | | | | | | | | | |
| 34 | **total** | 312 | | | | | | | | | | | | | | | | | | | | | | | | |
| 35 | | | | | | | | | | | | | | | | | | | | | | | | | | |